

ROBUST ENTANGLEMENT IN ATOMIC SYSTEMS VIA Λ -TYPE PROCESSES

Özgür Cakır, M. Ali Can, Alexander Klyachko, and Alexander Shumovsky
Faculty of Science, Bilkent University, Bilkent, Ankara, 06533 Turkey

It is shown that the system of two three-level atoms in Λ configuration in a cavity can evolve to a long-lived maximum entangled state if the Stokes photons vanish from the cavity by means of either leakage or damping. The difference in evolution picture corresponding to the general model and effective model with two-photon process in two-level system is discussed.

I. INTRODUCTION

During the last decade, the problem of engineered entanglement in atomic systems has attracted a great deal of interest (see [1,2,3,4,5,6,7,8] and references therein). In particular, the atomic entangled states were successfully realized through the use of cavity QED [1] and technique of ion traps [3]. At present, one of the most important problems under consideration is how to make a long-lived and easy-monitored atomic entangled state with existing experimental technique.

An interesting scheme has been proposed recently [9]. In this scheme, the two identical atoms are placed into a cavity tuned to resonance with one of the dipole-allowed transitions. Initially both atoms are prepared in the ground state, while the cavity field consists of a single photon. It is easy to show that the atom-field interaction leads in this case to a maximum atomic entangled state such that the single excitation is shared between the two atoms with equal probability. It was proposed in [9] to consider the absence of photon leakage from a nonideal cavity as a signal that the atomic entangled state has been created. The scheme can also be generalized on the case of any even number of atoms $2n$, sharing n excitations. In this case, the atomic entangled states are represented by the so-called $SU(2)$ phase states [10].

Another interesting proposal is to use a strong coherent drive to provide the multipartite entanglement in a system of two-level atoms in a high-Q cavity [11]. This approach can be used to produce the atomic entanglement as well as that of atoms and cavity modes and even of different cavity modes.

In the schemes of Refs. [9,11], the lifetime of the entanglement is defined by the specific time scale of the dipole-allowed radiative processes in atoms. Unfortunately this lifetime is usually quite short [12].

Generally speaking, the lifetime of atomic entanglement is specified by the interaction of atoms with environment. For example, in the model of Ref. [10] the environment is represented by the vacuum field that causes emission of a photon getting out of the cavity.

The interaction with environment can also be used to create a long-lived entanglement in atomic systems. For example, the initially non-entangled system may evolve to an entangled state connected with the atomic states that cannot be depopulated by radiation decay. In this case, the lifetime of the entangled state is specified by the considerably long nonradiative processes. Possible realization is provided by the use of three-level Λ -type process instead of the two-level scheme of Refs. [9,11]. The process is illustrated by Fig. 1. Here the levels 1 and 2 are connected by the electric dipole transitions as well as the levels 2 and 3. In turn, the dipole transition between the levels 3 and 1 is forbidden because of the parity conservation [13]. The absorption of pumping photon by the transition $1 \leftrightarrow 2$ with further jump of the electron to the level 3 accompanied by emission of "Stokes" photon can be interpreted as a kind of Raman process in atomic system with emission of Stokes photon (see [14] and references therein).

It is clear that the atom excited to the level 3 can change the state either by absorption of the Stokes photon resonant with respect to the transition $3 \leftrightarrow 2$ or through a nonradiative decay. Assume now that the two identical Λ -type atoms are placed into a cavity of high quality with respect to the pumping photons resonant to the transition $1 \leftrightarrow 2$. We also assume that the Stokes photons created by the transition $2 \rightarrow 3$ either leave the cavity freely or are absorbed by the cavity walls. Then, the atom-field interaction may lead to creation of maximum entangled atomic state

$$\frac{1}{\sqrt{2}}(|3, 1\rangle + |1, 3\rangle), \quad (1)$$

whose lifetime is determined by the slow processes of nonradiative decay.

The above scheme has been proposed in Ref. [10] and briefly discussed in Ref. [15]. The main objective of present paper is to consider in details the evolution towards the long-lived atomic entangled state (1) and feasibility of the scheme with present experimental technique.

The paper is organized as follows. In Sec. II we discuss the model Hamiltonians that can be used to describe the process under consideration. Viz, we discuss the model of a one-photon three-level interaction and an effective model of two-photon process in two-level system. Then, in Sec. III, we examine the irreversible dynamics in a cavity with leakage of Stokes photons, leading to the state (1). We show that both models describe the exponential evolution to the state (1). At the same time, the effective model that corresponds to a rough time scale is unable to take into account the possible oscillations of population between the states 1 and 2. Let us stress that the monitoring of Stokes photons outside the cavity can be used to detect the atomic entangled state (1) in the case under consideration.

Another way of creation of state (1) through the use of a cavity with very low quality with respect to the Stokes photons is discussed in Sec. IV. Finally, in Sec. V, we discuss the possible realization of the scheme under consideration.

II. THE MODELS OF Λ -TYPE PROCESS

Assume that a system of N identical three-level atoms with Λ -type transitions shown in Fig. 1 interacts with the cavity mode close to resonance with $1 \leftrightarrow 2$ transition and with the Stokes radiation that can leave the cavity freely. Then, following [13,14], we can choose the model Hamiltonian in the following form

$$\begin{aligned} H &= H_0 + H_{int}, \\ H_0 &= \omega_P a_P^\dagger a_P + \sum_k \omega_{Sk} a_{Sk}^\dagger a_{Sk} \\ &+ \sum_f [\omega_{21} R_{22}(f) + \omega_{31} R_{33}(f)], \end{aligned} \quad (2)$$

$$H_{int} = \sum_f \lambda_P R_{21}(f) a_P + \sum_{f,k} \lambda_{Sk} R_{23}(f) a_{Sk} + H.c. \quad (3)$$

Here a_P denotes the photon operator of the cavity mode with frequency ω_P , a_{Sk} describes annihilation of Stokes photon with frequency ω_{Sk} , and ω_{21} and ω_{31} are the frequencies of the corresponding atomic levels with respect to the ground level 1. The operator

$$R_{ij}(f) = |i_f\rangle\langle j_f|$$

gives the transition from level j to level i , index f marks the number of atom. In Eq. (3), λ_P and λ_{Sk} are the coupling constants, specifying the dipole transitions $2 \leftrightarrow 1$ and $3 \leftrightarrow 2$, respectively. Summation over k in (3) implies that the Stokes photons do not feel the presence of cavity walls. This summation involves the modes distributed with a certain density in a narrow band near the atomic transition frequency

$$\omega_S \equiv \omega_{23} = \omega_{21} - \omega_{31}, \quad (4)$$

corresponding to the natural line breadth.

Apart from the total electron occupation number, the model (2), (3) has the two integrals of motion

$$\begin{aligned} N_P &= a_P^\dagger a_P + \sum_f \{R_{22}(f) + R_{33}(f)\} \\ N_S &= \sum_k a_{Sk}^\dagger a_{Sk} - \sum_f R_{33}(f). \end{aligned} \quad (5)$$

Consider the system of only two atoms. Assume that both atoms are prepared initially in the ground states 1, the cavity contains a single photon of frequency ω_P , and Stokes field is in the vacuum state. Then, because of the integrals of motion (5), the evolution of the system occurs in a single-excitation domain of the Hilbert space spanned by the vectors

$$\begin{cases} |\psi_1\rangle &= |1, 1\rangle \otimes |1_P\rangle \otimes |0_S\rangle \\ |\psi_2^{(\pm)}\rangle &= \frac{1}{\sqrt{2}}(|1, 2\rangle \pm |2, 1\rangle) \otimes |0_P\rangle \otimes |0_S\rangle \\ |\psi_{3k}^{(\pm)}\rangle &= \frac{1}{\sqrt{2}}(|1, 3\rangle \pm |3, 1\rangle) \otimes |0_P\rangle \otimes |1_{Sk}\rangle \end{cases} \quad (6)$$

By construction, the four states (6) labeled by the superscripts $\pm$ manifest the maximum entanglement. No one of the states (6) is the eigenstate of the Hamiltonian (2) and (3). At the same time, it is easily seen that the action of operator (3) cannot transform the states

$$\{|\psi_1\rangle, |\psi_2^{(+)}\rangle, |\psi_{3k}^{(+)}\rangle\} \quad (7)$$

into the states

$$\{|\psi_2^{(-)}\rangle, |\psi_{3k}^{(-)}\rangle\} \quad (8)$$

and vice versa. Thus, the evolution of the system from the initial state $|\psi_1\rangle$ occurs in the subspace spanned by the three vectors (7). In this case, the states (8) can be discarded from the basis (6).

Instead of the one-photon three-level model described by the Hamiltonian (2) and (3), an effective model of two-photon process can also be used under a certain condition [16,17]. Viz, if the cavity is tuned consistent with two-photon energy conservation, i.e.

$$E_3 - E_1 = \omega_1 - \omega_2,$$

we are left with only one detuning parameter

$$\Delta = E_1 - E_2 - \omega_1 = E_2 - E_3 - \omega_S.$$

Here E_i denotes the energy of corresponding atomic level. Then, it was shown in [16] that under the condition

$$\Delta \gg E_3 - E_1$$

the dynamics of the system is governed by the effective Hamiltonian of the form

$$\begin{aligned} H^{eff} = & \omega_P a_P^\dagger a_P + \omega_S a_S^\dagger a_S + \sum_f \omega_{31} R_{33}(f) \\ & + \sum_f \lambda [R_{31}(f) a_S^\dagger a_P + a_P^\dagger a_S R_{13}(f)], \end{aligned} \quad (9)$$

describing the two-level transition with simultaneous absorption of pumping photon and creation of Stokes photon. Here λ is an effective coupling constant.

III. DYNAMICS DESCRIBED BY THE HAMILTONIAN (2),(3)

Under the assumption that there are only two three-level Λ -type atoms in the cavity and that the system is initially prepared in the state $|\psi_1\rangle$ in (5), in view of the results of previous section we should choose the time-dependent wave function as follows

$$|\Psi(t)\rangle = C_1|\psi_1\rangle + C_2|\psi_2\rangle + \sum_k C_{3k}|\psi_{3k}\rangle, \quad (10)$$

$$C_1(0) = 1, \quad C_2(0) = 0, \quad \forall k \quad C_k(0) = 0, \quad (11)$$

using the reduced basis (7). Here for simplicity we use the notations $|\psi_2\rangle \equiv |\psi_2^{(+)}\rangle$ and $|\psi_{3k}\rangle \equiv |\psi_{3k}^{(+)}\rangle$. The time-dependent Schrödinger equation with the Hamiltonian (2) and (3) then leads to the following set of equations for the coefficients in (11)

$$\begin{cases} i\dot{C}_1 = \omega_P C_1 + \lambda_P \sqrt{2} C_2 \\ i\dot{C}_2 = \omega_{21} C_2 + \lambda_P \sqrt{2} C_1 + \sum_k \lambda_{Sk} C_{3k} \\ i\dot{C}_{3k} = (\omega_{31} + \omega_{Sk}) C_{3k} + \lambda_{Sk} C_2 \end{cases} \quad (12)$$

To find solutions of (12), let us represent the last equation in (12) in the form

$$C_{3k}(t) = -i\lambda_{Sk} \int_0^t C_2(\tau) e^{i(\omega_{31} + \omega_{Sk})(\tau - t)} d\tau. \quad (13)$$

Then, we should take the time derivative in both sides of the first equation in (12) and substitute the second equation together with integral representation (13). We get

$$\begin{aligned} i\ddot{C}_1 = & (\omega_P + \omega_{21})\dot{C}_1 + i(\omega_{21}\omega_P - 2\lambda_P^2)C_1 \\ & - \sum_k \lambda_{Sk}^2 \int_0^t (i\dot{C}_1 - \omega_P C_1) e^{i(\omega_{31} + \omega_{Sk})(\tau - t)} d\tau. \end{aligned}$$

Carrying out the integration by parts, we get the following integro-differential equation with respect to only one unknown variable $C_1(t)$:

$$\begin{aligned}
i\ddot{C}_1 = & (\omega_P + \omega_{21})\dot{C}_1 + i(\omega_{21}\omega_P - 2\lambda_P^2 \\
& - \sum_k \lambda_{Sk}^2)C_1 + i \sum_k \lambda_{Sk}^2 e^{-i(\omega_{31} + \omega_{Sk})t} - \sum_k \lambda_{Sk}^2 (\omega_{31} + \omega_{Sk} \\
& - \omega_P) \int_0^t C_1(\tau) e^{i(\omega_{31} + \omega_{Sk})(\tau-t)} d\tau
\end{aligned} \tag{14}$$

This integro-differential equation can be analyzed through the use of Laplace transformation, which is usually considered in the context of natural line breadth (see Ref. [19]). Let us emphasize that, in contrast to the conventional Wigner-Weisskopf theory, Eq. (14) contains the second-order derivative in addition to the first-order term. In other words, it should describe oscillations together with the decay. We have

$$\begin{aligned}
\int_0^\infty C_1(t) e^{-st} dt &= \mathcal{L}(C_1), \\
\int_0^\infty \dot{C}_1(t) e^{-st} dt &= s\mathcal{L} - 1, \\
\int_0^\infty \ddot{C}_1(t) e^{-st} dt &= s^2\mathcal{L} - s - \dot{C}_1(0) = s^2\mathcal{L} - s + i\omega_P.
\end{aligned}$$

Then, Eq. (14) is reduced into the following algebraic equation with respect to $\mathcal{L}$:

$$\begin{aligned}
& \mathcal{L}[is^2 - s(\omega_P + \omega_{21}) - i(\omega_{21}\omega_P - 2\lambda_P^2 - \sum_k \lambda_{Sk}^2)] \\
&= -(\omega_P + \omega_{21}) + i \sum_k \lambda_{Sk}^2 \frac{s + i(\omega_{31} + \omega_{Sk})}{s^2 + (\omega_{31} + \omega_{Sk})^2} - \int_0^\infty e^{-st} \left\{ \sum_k \lambda_{Sk}^2 \right. \\
& \quad \times (\omega_{31} + \omega_{Sk} - \omega_P) \int_0^t C_1(\tau) e^{i(\omega_{31} + \omega_{Sk})(\tau-t)} d\tau \Big\} dt.
\end{aligned} \tag{15}$$

The last term in the right-hand side of this expression can be represented as follows

$$\begin{aligned}
& \int_0^\infty e^{-st} \left\{ \int_0^t e^{i(\omega_{31} + \omega_{Sk})(\tau-t)} C_1(\tau) d\tau \right\} dt \\
&= \int_0^\infty C_1(\tau) e^{i(\omega_{31} + \omega_{Sk})\tau} d\tau \int_\tau^\infty e^{-[s + i(\omega_{31} + \omega_{Sk})]t} dt \\
&= \frac{\mathcal{L}}{f + i(\omega_{31} + \omega_{Sk})}.
\end{aligned}$$

Thus, Eq. (15) takes the form

$$\begin{aligned}
\mathcal{L} = & \left[i \sum_k \lambda_{Sk}^2 \frac{s + i(\omega_{31} + \omega_{Sk})}{s^2 + (\omega_{31} + \omega_{Sk})^2} - (\omega_P + \omega_{21}) \right] \\
& \times [is^2 - s(\omega_P + \omega_{21}) + i(\omega_P\omega_{21} - 2\lambda_P^2 - \sum_k \lambda_{Sk}^2) \\
& + \sum_k \frac{\lambda_{Sk}^2 (\omega_{31} + \omega_{Sk} - \omega_P)}{s + i(\omega_{31} + \omega_{Sk})}]^{-1}.
\end{aligned} \tag{16}$$

Then, the exact form of the time behavior of the coefficient $C_1(t)$ in Eq. (10) is governed by the inverse Laplace transformation

$$C_1(t) = \frac{1}{2\pi i} \int_{\epsilon-i\infty}^{\epsilon+i\infty} e^{st} \mathcal{L}(s) ds, \tag{17}$$

where ϵ is a small real positive number and s in the integrand is considered as a complex argument. As soon as the explicit time behavior of $C_1(t)$ is known, the other coefficients in (10) can be defined through the use of Eqs. (12) and (13).

In particular, it follows from Eqs. (6), (10), and (11) that the probability to have the atomic entangled state (1) has the form

$$\begin{aligned} \sum_k |C_{3k}|^2 &= 1 - |C_1(t)|^2 - |C_2(t)|^2 \\ &= 1 - |C_1(t)|^2 - \frac{|i\dot{C}_1(t) - \omega_P C_1(t)|^2}{2\lambda_P^2}. \end{aligned} \quad (18)$$

Thus, the Eqs. (16) and (17) completely determine the probability to have the robust entangled state (1).

It can be shown that Eq. (16) describes the reversible, Poincaré type behavior (e.g., see [19]). The irreversible evolution can be obtained under the further assumption that the atomic transition $2 \leftrightarrow 3$ interacts with continuum of Stokes modes rather than with a discrete spectrum. Similar assumption is used within conventional Wigner-Weisskopf theory [18]. This means that we replace summation over k by integration over $\omega = ck$:

$$\sum_k \cdots \rightarrow \int_{-\infty}^{\infty} \cdots \rho(\omega) d\omega.$$

Here the measure $\rho(\omega)d\omega$ defines the density of states of Stokes photons with different frequencies corresponding to the natural line breadth. Unlike conventional Wigner-Weisskopf theory, Eqs. (16) and (17) describe a superposition of exponential decay and harmonic oscillations caused by the interaction between the $1 \leftrightarrow 2$ transitions and cavity field.

Further analysis shows that, the coefficients C_1 and C_2 up to the second order in $\lambda/(\Gamma - i\Delta_P)$ are

$$\begin{aligned} C_1(t) &\approx \left[-\frac{2\lambda^2}{(\Gamma - i\Delta)^2} e^{(-\Gamma + i\Delta)t} + \left(1 + \frac{2\lambda^2}{(\Gamma - i\Delta)^2}\right) e^{-\frac{2\lambda^2}{\Gamma - i\Delta}t} \right] e^{-i\omega_P t} \\ C_2(t) &\approx -\frac{\sqrt{2}\lambda}{i\Gamma + \Delta} \left[e^{-\Gamma t} - e^{-(\frac{2\lambda^2}{\Gamma - i\Delta} + i\Delta)t} \right] e^{-i\omega_{21}t} \end{aligned} \quad (19)$$

where

$$\Delta_P = \omega_P - \omega_{12}$$

is the detuning factor for the pumping mode and

$$\Gamma = \rho(\omega_S) \lambda_{Sk} (k = \omega_S/c), \quad \omega_S = \omega_{21} - \omega_{31}.$$

Since $\rho(\omega_S) \gg 1$, $\Gamma \gg \lambda_P, \lambda_{Sk}$ and the result (19) corresponds to the second order order with respect to $\lambda_P/(\Gamma - i\Delta)$.

It is seen that Eq. (19) describes the damped oscillations of the coefficient $C_1(t)$ in (10). According to (11), coefficient $C_2(t)$ manifests similar behavior. Thus, the probability (18) to get the robust entangled state tends to unit as $t \rightarrow \infty$ (see Fig. 2). The decay time γ^{-1} is defined by the coupling constant and detuning parameter for the pumping mode and by the width of the Stokes line specified by the parameter Γ . Although Eq. (19) contains oscillating terms, the decay is quite strong and the influence of oscillating terms cannot be observed even when $\Delta_P \leq \Gamma$. Only when detuning strongly exceeds Γ , the oscillations start to contribute significantly into the time evolution of probability (18).

Similar result can also be obtained in terms of the effective Hamiltonian (9) [15]. It should be stressed that the assumptions made in the process of derivation of (9) lead to an effective "roughing" of the time scale. The obliteration of the level 2 leads to the simultaneous neglect of the Rabi oscillations between the levels 1 and 2. Therefore, the effective model (9) is incapable of recognition of time oscillations and gives only rough picture of purely exponential evolution.

While the atomic system evolves to the maximum entangled state (1), the Stokes photon leaves the cavity. Thus, the observation of Stokes photon outside the cavity can be considered as a signal that the robust entangled state has been prepared.

IV. CAVITY WITH ABSORPTION OF STOKES PHOTONS

The atomic entangled state (1) can also be realized when the Stokes mode is strongly damped in the cavity. For simplicity, we still assume no damping for the pumping mode. At the same time, the Stokes photons are supposed to

be absorbed by the cavity walls. In other words, we consider the case of high quality (with respect to pumping field photons) and zero-temperature cavity, corresponding to the most of the Ridberg maser experiments [20,21]. In this case, the effect of damping can be calculated through the use of the so-called dressed-atom approximation [22].

The model Hamiltonian, describing the process under consideration, can be chosen as follows

$$\begin{aligned}
H &= H_0 + H_{int}, \\
H_0 &= \omega_P a_P^\dagger a_P + \omega_S a_S^\dagger a_S \\
&+ \sum_f [\omega_{21} R_{22}(f) + \omega_{31} R_{33}(f)], \\
H_{int} &= \sum_f [\lambda_P R_{21}(f) a_P + \lambda_S R_{23}(f) a_S] + H.c.
\end{aligned} \tag{20}$$

This corresponds to the single-Stokes-mode approximation in (2) and (3). The eigenstates of the Hamiltonian (20) have the form

$$\begin{aligned}
|\psi_0\rangle &= \frac{\lambda_S}{\epsilon} |\psi_1\rangle - \frac{\lambda_P \sqrt{2}}{\epsilon} |\psi_3\rangle, \\
|\psi_\pm\rangle &= \pm \frac{\lambda_P}{\epsilon} |\psi_1\rangle + \frac{1}{\sqrt{2}} |\psi_2\rangle \pm \frac{\lambda_S}{\epsilon \sqrt{2}} |\psi_3\rangle
\end{aligned} \tag{21}$$

where $|\psi_1\rangle$ coincides with the first state in (6), $|\psi_2\rangle = |\psi_2^{(+)}\rangle$, and

$$|\psi_3\rangle = \frac{1}{\sqrt{2}} (|3, 1\rangle + |1, 3\rangle) \otimes |0_P\rangle \otimes |1_S\rangle.$$

In Eqs. (21),

$$\epsilon = \sqrt{2\lambda_P^2 + \lambda_S^2}.$$

Under the assumption of exact resonance

$$\omega_P = \omega_{21} = \omega_{31} + \omega_S$$

that we use hereafter for simplicity, the corresponding eigenvalues are

$$H|\psi_0\rangle = \omega_P |\psi_0\rangle, \quad H|\psi_\pm\rangle = (\omega_P \pm \epsilon) |\psi_\pm\rangle.$$

Besides that, there is one more eigenstate

$$|\psi_4\rangle = \frac{1}{\sqrt{2}} (|3, 1\rangle + |1, 3\rangle) \otimes |0_P\rangle \otimes |0_S\rangle, \tag{22}$$

such that

$$H|\psi_4\rangle = \omega_{31} |\psi_4\rangle.$$

It is clear that this eigenstate corresponds to the maximum atomic entanglement (1). Physically, this state is achieved when the Stokes photon is absorbed by the cavity walls.

To take into account the cavity damping of Stokes photons, consider interaction with a "phonon reservoir" responsible for the absorption of photons by cavity walls [18]. Then, the Hamiltonian (20) should be supplemented with the term

$$H_{loss} = \sum_q \eta_q (b_q^\dagger a_S + a_S^\dagger b_q) + \sum_q \Omega_q b_q^\dagger b_q, \tag{23}$$

where $b_q, b_q^\dagger$ are the Bose operators of "phonons" in the cavity walls.

The density matrix of the system can be chosen as follows

$$\rho(t) = \sum_{j,\ell} \rho_{j\ell}(t) |\psi_j\rangle \langle \psi_\ell|, \quad j, \ell = 0, \pm, 4, \tag{24}$$

where $|\psi_j\rangle$ are the eigenstates (21) and (22) and $\rho_{j\ell}(t)$ denote the time-dependent c -number coefficients.

With the total Hamiltonian

$$H_{tot} = H + H_{loss}$$

in hand, we can now write the Master Equation, eliminating the cavity degrees of freedom (e.g., see [23]),

$$\dot{\rho} = -i[H, \rho] + \kappa\{2a_S\rho a_S^\dagger - a_S^\dagger a_S\rho - \rho a_S^\dagger a_S\}, \quad (25)$$

so that the contribution of (23) is taken into account effectively through the Liouville term. Here $1/\kappa$ is the lifetime of a Stokes photon in the cavity and $Q = E_3/\kappa$ is the quality factor with respect to the Stokes photons (E_3 denotes the energy of the level 3). Let us choose the same initial condition as in previous section, so that

$$\rho(0) = |\psi_1\rangle\langle\psi_1| \quad (26)$$

where the initial state $|\psi_1\rangle$ is expressed in terms of the eigenstates (21) as follows

$$|\psi_1\rangle = \frac{\lambda_S}{\epsilon}|\psi_0\rangle + \frac{\lambda_P}{\epsilon}(|\psi_+\rangle - |\psi_-\rangle).$$

The equation (25) can now be solved numerically at different values of parameter κ , specifying the absorption of Stokes photons. The results are shown in Fig. 3. It is seen that the system evolves to the robust atomic entangled state (1). The stairs-like structure is again caused by competition between the transitions $1 \leftrightarrow 2$ and $2 \leftrightarrow 3$. Although such a behavior is an inherent property of the model under consideration, the stairs become more visible with increase of κ (see Fig. 3).

Similar result can also be obtained within the framework of effective model with the Hamiltonian (9) and damping described by Eq. (23). In this case, the density matrix consists of only six elements because the state $|\psi_0\rangle$ in (21) should be discarded and the states $|\psi_\pm\rangle$ are changed by the states

$$|\phi_\pm\rangle = \frac{1}{\sqrt{2}}(|\psi_1\rangle \pm |\psi_3\rangle)$$

with the eigenvalues

$$\varepsilon_\pm = \omega_P \pm \lambda\sqrt{2}.$$

It should be stressed that the effective model does not show the stairs-like behavior of $\rho_{44}(t)$.

V. SUMMARY AND DISCUSSION

In this paper, we have studied the quantum dynamics of a system of two three-level atoms in the Λ configuration interacting with two modes of quantized electromagnetic field in a cavity under the assumption that the Stokes-mode photons either leave the cavity freely or are damped rapidly. It is shown that in both cases the system evolves from the state when both atoms are in the ground state and cavity contains a pumping photon into the robust entangled state (1). The lifetime of this final state is defined completely by the nonradiative processes and is therefore relatively long.

In the case of cavity transparent for the Stokes photons, the creation of Stokes photon signals the rise of atomic entanglement. Such a photon can be monitored outside the cavity.

Let us stress that the general models with the Hamiltonians (2), (3) and (20), that takes into account all three atomic levels, admit a certain peculiarities in the evolution towards the robust entangled state caused by the competition of transitions $1 \leftrightarrow 2$ and $2 \leftrightarrow 3$. The effective model with adiabatically eliminated highest excited level is incapable of description of these peculiarities, while predicts correct asymptotic behavior. Moreover, the general model admits also a number of intermediate maximum entangled states ($|\psi_2\rangle$ and $|\psi_{3k}\rangle$ in Eq. (6)) that do not exist in the effective model. Unfortunately, the lifetime of these entangled states are defined by the dipole radiative processes and are therefore too short.

It seems to be possible to realize the robust entangled state (1) using the present experimental technique. Let us note in this connection that the single-atom Raman process has been recently observed [24].

We also note that the way of experimental preparation of single-photon cavity excitation is well known [21]. The atoms can propagate through the cavity, using either the same opening or two different openings. All measurements aimed at the detection of atomic entanglement can be performed outside the cavity.

Although our results were obtained for a system of two atoms, they can be generalized with easy on the case of big atomic clusters, using the method of Ref. [10]. In fact, it is possible to show that a certain robust entanglement can be obtained in a system with any even number $2N$ of three-level Λ -type atoms initially prepared in the ground state and interacting with N pumping photons.

ACKNOWLEDGEMENT



One of the authors (A.Sh.) would like to thank Prof. J.H. Eberly, Prof. A. Vourdas, and Prof. D.G. Welsch for many useful discussions.


-
- [1] E. Hagley, X. Maitre, G. Nogues, C. Wunderlich, M. Brune, J.M. Raymond, and S. Haroche, *Phys. Rev. Lett.* **79**, 1 (1997).
 - [2] A. Furusawa, J.L. Sørensen, S.L. Braunstein, C.A. Fuchs, H.J. Kimble, and E.S. Polzik, *Science* **282**, 706 (1998).
 - [3] Q.A. Turchette, C.S. Wood, B.E. King, C.J. Myatt, D. Leibfried, W.M. Itano, C. Monroe, and D.J. Wineland, *Phys. Rev. Lett.* **81**, 3631 (1998).
 - [4] B. Bederson and H. Walther, editors, *Advances in Atomic, Molecular, and Optical Physics*, Vol. **42** (Academic Press, New York, 2000).
 - [5] C.J. Myatt, B.E. King, Q.A. Turchette, C.A. Sackett, D. Kielpinski, W.H. Itano, C. Monroe, and D.J. Wineland, *Nature*, **403**, 269 (2000).
 - [6] G. Rempe, *Ann. Phys. (Leipzig)* **9**, 843, (2000).
 - [7] J.M. Raymond, M. Brune, and S. Haroche, *Rev. Mod. Phys.* **73**, 565 (2001).
 - [8] B. Julsgaard, A. Kozhekin, and E. Polzik, *Nature* **413**, 400 (2001).
 - [9] M.B. Plenio, S.F. Huelga, A. Beige, and P.L. Knight, *Phys. Rev. A* **59**, 2468 (1999); A. Beige, S. Bose, D. Braun, S.F. Huelga, P.L. Knight, M.B. Plenio, and V. Verdál, *J. Mod. Optics* **47**, 2583 (2000).
 - [10] M.A. Can, A.A. Klyachko, and A.S. Shumovsky, *Phys. Rev. A* **66**, 022111 (2002).
 - [11] E. Solano, G.S. Agarwal, and H. Walther, quant-ph/0202071.
 - [12] E.U. Condon and G.H. Shortley, *The Theory of Atomic Spectra* (Cambridge University Press, New York, 1987); C.E. Moore, *Atomic Energy Levels* (US National Bureau of Standards, Washington DC, 1971).
 - [13] H.I. Yoo and J.H. Eberly, *Phys. Rep.* **118**, 239 (1985); F.L. Kien and A.S. Shumovsky, *Int. J. Mod. Phys B* **5**, 2287 (1991); R. Shore and P.L. Knight, *J. Mod. Optics* **40**, 1195 (1993).
 - [14] T. Quang and A.S. Shumovsky, in *Interaction of Electromagnetic Field with Condensed Matter*, edited by N.N. Bogolubov, A.S. Shumovsky, and V.I. Yukalov (World Scientific, Singapore, 1990).
 - [15] M.A. Can, A.A. Klyachko, and A.S. Shumovsky, quant-ph/0207006 (to be published in *Appl. Phys. Lett.* **81** No 20 (2002)).
 - [16] C.C. Gerry and J.H. Eberly, *Phys. Rev. A* **42**, 6805 (1990).
 - [17] C.K. Law and J.H. Eberly, *Phys. Rev. A* **47**, 3195 (1993); R.R. Puri, C.K. Law, and J.H. Eberly, *Phys. Rev. A* **50**, 4212 (1994).
 - [18] W.H. Louisell, *Radiation and Noise in Quantum Electronics* (Mc-Grow Hill, New York, 1964); C.W. Gardiner and P. Zoller, *Quantum Noise* (Springer, Berlin, 2000).
 - [19] R. Omnes, *Interpretation of Quantum Mechanics* (Princeton Univ. Press, NJ, 1994).
 - [20] G. Rempe, H. Walther, and N. Klein, *Phys. Rev. Lett.* **58**, 353 (1987); G. Rempe, F. Schmidt-Kaler, and H. Walther, *Phys. Rev. Lett.* **64**, 2783 (1990).
 - [21] H. Walther, in *Quantum Optics and the Spectroscopy of Solids*, edited by T. Hakio'uglu and A.S. Shumovsky (Kluwer, Dordrecht, 1997); B.T.H. Varcoe, S. Bratke, B.-G. Englert, and H. Walther, *Laser Physics*, **10**, 1 (2000); H. Walther, in *Quantum Communication, Computing, and Measurement*, **3**, edited by P. Tombesi and O. Hirota (Kluwer Academic/Plenum Press, New York, 2001).
 - [22] S. Haroche and J.M. Raymond, in *Advances in Atomic and Molecular Physics*, Vol. 20, edited by D.R. Bates and B. Bederson (Academic Press, New York, 1985); S.M. Barnett and P.L. Knight, *Phys. Rev. A* **33**, 2444 (1986); R.R. Puri and G.S. Agarwal, *Phys. Rev. A* **35**, 3433 (1987); H.T. Dung and A.S. Shumovsky, *Phys. Lett. A* **169**, 379 (1992).
 - [23] S.M. Barnett, P. Radmore *Methods in Theoretical Quantum Optics* (Clarendon Press, Oxford, 1997).
 - [24] M. Hennrich, T. Legero, A. Kuhn, and G. Rempe, *Phys. Rev. Lett.* **85**, 4872 (2000).


FIG. 1. Scheme of the process and configuration of atomic levels and transitions.

FIG. 2. Time evolution of probability (18) to have the robust entanglement at $\lambda_P = 0.001\Gamma$ for (1) $\Delta_P = 0$ (2) $\Delta_P = \Gamma$ (3) $\Delta_P = 2\Gamma$ (4) $\Delta_P = 4\Gamma$


FIG. 3. Time evolution of $\rho_{44}(t)$ at $\lambda_P = \lambda_S$ and $\kappa = 0.01\lambda_P$ (a) and $\kappa = 0.1\lambda_P$ (b).







(a)



(b)